High-Precision Penning-Trap Mass Spectrometry of Actinides at TRIGA-Trap

Tanvir Sayed 1Szilard Nagy 1, †Klaus Blaum 1Michael Block 2,3,4Burcu Cakirli 1Stanislav Chenmarev 1Christoph Düllmann 2,3,4Dennis Renisch 2,3

¹Max-Planck-Institut für Kernphysik, 69117 Heidelberg, Germany
²Helmholtz-Institut Mainz, 55128 Mainz, Germany
³Department Chemie - Standort TRIGA, Johannes Gutenberg-Universität Mainz, 55128 Mainz, Germany
⁴GSI Helmholtzzentrum für Schwerionenforschung, 64291 Darmstadt, Germany
[†]corresponding author's email: szilard.nagy@mpi-hd.mpg.de

Penning traps are widely used in high-precision mass spectrometry to determine atomic masses with exceptional precision and accuracy, playing a crucial role in atomic and nuclear physics research [1]. TRIGA-Trap is a high-precision, double penning-trap mass spectrometer located in the reactor hall of the TRIGA (Training, Research, Isotopes, General Atomic) research reactor in Mainz, Germany [2]. It also serves as one of the development platforms for the planned MATS (Measurements of very short-lived nuclides using an Advanced Trapping System) experiment at FAIR (Facility for Antiproton and Ion Research) which aims to investigate nuclei far from stability in order to enhance our knowledge on their fundamental nuclear properties and interactions [3, 4]. While the construction of FAIR is still underway, development platforms such as TRIGA-Trap conduct studies to optimise future experiments and test new emerging ideas.

At TRIGA-Trap, mass measurements of radioactive nuclides – particularly actinides – are performed with the PI-ICR (Phase-Imaging Ion-Cyclotron Resonance) technique. This method offers high sensitivity, resolving power and accuracy, while requiring relatively short measurement times [2]. Recent mass measurements of actinides, including ²⁴⁴Pu, ²⁴¹Am, ²⁴³Am, ²⁴⁸Cm, and ²⁴⁹Cf have achieved uncertainties at the parts-per-billion (ppb) level [5]. Currently, mass measurements in the Pu isotopic chain – including ²³⁸Pu, ²³⁹Pu, ²⁴⁰Pu, and ²⁴²Pu – are planned and in progress. These nuclides are in the vicinity of the neutron number N = 152, a region associated with a deformed sub-shell closure. The precise mass measurements allow the exploration of nuclear structure through trends in mass filters, such as S_{2n} (two-neutron separation energies) and $\delta V_{p,n}$ (average *p-n* interaction of the most loosely-bound two nucleons), as well as their differentials [5].

This presentation will provide an overview of the current status of the experiment, including recent mass measurements, their application in nuclear structure evaluation, and the investigations into systematic uncertainties that contribute to accurate mass determinations.

References

[1] J. Dilling, K. Blaum, M. Brodeur et al, Annu. Rev. Nucl. Part. Sci. 68(1) (2018).

- [2] S. Chenmarev, S. Nagy, J.J.W. van de Laar et al, Eur. Phys. J. A 59(2), 29 (2023).
- [3] D. Rodríguez, K. Blaum, W. Nörtershäuser et al, Eur. Phys. J. Spec. Top. 183(1), 1-123 (2010).
- [4] G. Rosner, Nucl. Phys. B Proc. Suppl. 167, 77–81 (2007).
- [5] S. Chenmarev, K. Blaum, M. Block et al, Eur. Phys. J. A 60, 204 (2024).